

After all potentially impairing conditions have been identified and the correct ratings recorded, the evaluator should select the clinically most appropriate (ie, most specific) method(s) and record the estimated impairment for each. The **cross-usage chart** (Table 17-2) indicates which methods and resulting impairment ratings may be combined. It is the responsibility

of the evaluating physician to explain in writing why a particular method(s) to assign the impairment rating was chosen. When uncertain about which method to choose, the evaluator should calculate the impairment using different alternatives and choose the method or combination of methods that gives the most clinically accurate impairment rating.

**Table 17-2** Guide to the Appropriate Combination of Evaluation Methods

Open boxes indicate impairment ratings derived from these methods can be combined.

	Limb Length Discrepancy	Gait Derangement	Muscle Atrophy	Muscle Strength	ROM Ankylosis	Arthritis (DJD)	Amputation	Diagnosis-Based Estimates (DBE)	Skin Loss	Peripheral Nerve Injury	Complex Regional Pain Syndrome (CRPS)	Vascular
Limb Length Discrepancy		X					X					
Gait Derangement	X		X	X	X	X	X	X	X	X	X	X
Muscle Atrophy		X		X	X	X	X	X		X	X	
Muscle Strength		X	X		X	X		X	*	X	0	
ROM Ankylosis		X	X	X		X		X			0	
Arthritis (DJD)		X	X	X	X							
Amputation	X	X	X	X								
Diagnosis-Based Estimates (DBE)		X	X	X	X							
Skin Loss		X										
Peripheral Nerve Injury		X	X	X							X	
Complex Regional Pain Syndrome (CRPS)		X	X	0	0					X		X
Vascular		X									X	

X = Do not use these methods together for evaluating a single impairment.

0 = See specific instructions for CRPS of the lower extremity.